

## Hydrogen Peroxide Kills *Staphylococcus aureus* by Reacting with Staphylococcal Iron to Form Hydroxyl Radical\*

(Received for publication, May 1, 1981)

John E. Repine‡, Richard B. Fox§, and Elaine M. Berger

From the Webb-Waring Lung Institute and the Pulmonary Divisions of the Departments of Medicine and Pediatrics of the University of Colorado Health Sciences Center, Denver, Colorado 80262

Two lines of investigation supported the premise that killing of *Staphylococcus aureus*, 502A, by hydrogen peroxide involves formation of the more toxic hydroxyl radical ( $\cdot\text{OH}$ ) through the iron-dependent Fenton reaction. First, growing *S. aureus* overnight in broth media with increasing concentrations of iron increased their content of iron and dramatically enhanced their subsequent susceptibility to killing by  $\text{H}_2\text{O}_2$ . Second, in direct relation to their effectiveness as  $\cdot\text{OH}$  scavengers, thiourea, dimethyl thiourea, sodium benzoate, and dimethyl sulfoxide inhibited  $\text{H}_2\text{O}_2$ -mediated killing of *S. aureus*.

The crucial role of iron in the resolution between microbial invader and host remains unsettled (1). A majority of the evidence indicates that hypoferremia is mainly beneficial to the host. This conclusion is largely based on knowledge that iron is an essential growth requirement for microbial organisms but is also supported by observations suggesting an increased propensity to infection during hyperferremic episodes, such as those occurring during liver destruction, hemolytic anemia, and other disorders. Support for the possibility that iron helps the host relates primarily to findings that severe hypoferremia may impair host defense mechanisms, such as the killing of microorganisms by phagocytic cells (2).

To further investigate the possible biochemical role of iron in bacterial host defense interactions, we recently grew bacteria in iron before incubating them with neutrophils (3). We found that while preincubating *Staphylococcus aureus* in iron did not increase their susceptibility to killing by neutrophils, it markedly increased their susceptibility to killing by hydrogen peroxide (3).


In the present investigation we tried to elucidate the mech-

\* This work was supported in part by Grants HL24248 and BRSG-RR-05357 from the National Institutes of Health, the American Heart Association, The Council for Tobacco Research-U.S.A., Inc., and The Kroc Foundation. Presented in part at the Gordon Conference on Oxygen Radicals in Biology and Medicine, Ventura, California, 1981. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ Established Investigator of the American Heart Association. To whom reprint requests and correspondence should be sent at Webb-Waring Lung Institute, University of Colorado Health Sciences Center, 4200 East Ninth Ave., C-322, Denver, Colorado 80262.

§ Clinical Fellow of the National Institutes of Health.

anism underlying this finding by addressing the hypothesis that iron in microorganisms facilitates bactericidal mechanisms by reacting with  $\text{H}_2\text{O}_2$  to form the more toxic hydroxyl radical ( $\cdot\text{OH}$ ). Production of hydroxyl radical occurs by the well known Fenton reaction (4-7),



The results of the present investigation supported this premise. Increasing the concentration of iron in *S. aureus* increased their susceptibility to killing by  $\text{H}_2\text{O}_2$ . Scavengers of  $\cdot\text{OH}$  inhibited killing of *S. aureus* by  $\text{H}_2\text{O}_2$  and in direct relation to their effectiveness as  $\cdot\text{OH}$  scavengers. This mechanism may be a new way in which iron is beneficial to the host.

### EXPERIMENTAL PROCEDURES

*S. aureus*, 502A, were initially grown overnight (18 h) at 37 °C in Bacto nutrient broth media with increasing concentrations of  $\text{FeSO}_4$ , harvested, washed thoroughly, and their iron content (micrograms/ $10^9$  organisms) determined using the ferozine method (8). Mean iron concentrations of the overnight broth media of 0.3, 3.3, or 6.5  $\mu\text{g}/\text{ml}$  yielded *S. aureus* with mean iron concentrations of 0.2, 0.7, or 1.7  $\mu\text{g}/10^9$  organisms, respectively. Each iron value is the mean of 15 to 20 determinations. Killing was measured by mixing *S. aureus* ( $5 \times 10^6$ ) in human serum albumin (0.1%) and Hanks' balanced salt solution (1 ml) with or without  $\text{H}_2\text{O}_2$  (9). After mixing, each tube was immediately sampled by micropipette, capped, and incubated at 37 °C. After 60 min, each tube was resampled. Samples were diluted and plated on agar. After 24 h, the number of colonies on each plate was counted electronically (Automatic Colony Counter, 3M, Minneapolis, MN). The percentage of the initial inoculum of *S. aureus* killed was calculated from 0- and 60-min plate counts and expressed as the mean  $\pm$  S.E. Each  $\text{LD}_{50}$  point on the graph is the average  $\pm$  S.E. of 12 determinations.

Methane production from  $\text{Me}_2\text{SO}$ <sup>1</sup> was measured by gas chromatography as previously described (10). In some samples,  $\gamma$ -irradiation was administered for 2 min at 0 °C in closed glass vials in a  $\gamma$  cell <sup>60</sup>Co source delivering a dose of roughly 1.8 krad/min (18  $\gamma$ -rays/min) as described (11).  $\text{CH}_4$  production from  $\text{Me}_2\text{SO}$  which has been irradiated is a relatively specific consequence of reaction of  $\text{Me}_2\text{SO}$  with  $\cdot\text{OH}$  (10, 11) and, therefore, is an acceptable way of assessing the relative  $\cdot\text{OH}$  scavenging abilities of these inhibitors under our test conditions.

### RESULTS

Two lines of investigation supported the premise that killing of *S. aureus* by  $\text{H}_2\text{O}_2$  involves production of the more toxic hydroxyl radical by an iron-dependent Fenton mechanism. First, preincubation of *S. aureus*, 502A, overnight (18 h) in broth media with increasing concentrations of  $\text{FeSO}_4$  progressively increased their intrinsic content of iron and greatly enhanced their subsequent susceptibility to killing by  $\text{H}_2\text{O}_2$  in the absence of extrinsic iron added during the bactericidal assay (Fig. 1). In contrast, addition of iron (0.3 to 6.5  $\mu\text{g}/\text{ml}$ ) during the bactericidal assay did not appreciably increase killing by  $\text{H}_2\text{O}_2$  of *S. aureus* initially grown overnight in broth media without added iron (data not shown). Thus, with respect to susceptibility to  $\text{H}_2\text{O}_2$ , the intrinsic iron concentrations of the *S. aureus* are much more important than the amount of iron present extrinsically.

The suggested role of iron was further supported by control studies showing that direct alterations in the growth rates, viability, and catalase activities of *S. aureus* grown in increasing concentrations of iron did not account for their amplified sensitivity to killing by  $\text{H}_2\text{O}_2$ . Specifically, adding  $\text{FeSO}_4$  to

<sup>1</sup> The abbreviation used is:  $\text{Me}_2\text{SO}$ , dimethyl sulfoxide.




FIG. 1. Hydrogen peroxide needed to kill 50% of the initial inoculum of *S. aureus* ( $LD_{50}$ ) as a function of the intrinsic iron (micrograms/ $10^9$  organisms) of the *S. aureus* (○—○). Increasing the iron content of the *S. aureus* markedly increased their susceptibility to killing by  $H_2O_2$ . Each  $LD_{50}$  point on the graph is the mean  $\pm$  S.E. of 12 determinations.

their overnight broth media did not change the growth rates or viability of *S. aureus* during the bactericidal assay. The numbers of *S. aureus* in 10- $\mu$ l samples taken from *S. aureus* grown in broth mixtures with 0.3 or 6.5  $\mu$ g/ml of iron were the same after 15 (2338  $\pm$  42 versus 2267  $\pm$  74), 17 (2194  $\pm$  42 versus 2201  $\pm$  29), 19 (2210  $\pm$  125 versus 1925  $\pm$  126), or 20.5 (1954  $\pm$  67 versus 1875  $\pm$  53) h of incubation (6 determinations for each value). In the absence of  $H_2O_2$ , the viable numbers of *S. aureus* at the end of the 1-h killing assay were 1.05  $\pm$  0.09 times the numbers at the beginning of the bactericidal assay. This was true whether *S. aureus* had been previously grown overnight in media containing either 0.3 or 6.5  $\mu$ g/ml of iron (12 determinations). In addition, preliminary growth in broth media with 0.3 or 6.5  $\mu$ g/ml of iron did not change the catalase activities of *S. aureus* which were, respectively, 2.6  $\pm$  0.2 or 2.8  $\pm$  0.2  $\mu$ M  $H_2O_2$  decomposed/ $10^9$  *S. aureus*/min (5 determinations) (12). Furthermore, since no peroxidase activity was found in either high iron or low iron *S. aureus*, it is unlikely that the observed alterations in killing involved Fe-mediated alterations in peroxidase (13).

The second line of evidence implicating participation of  $\cdot$ OH in killing of *S. aureus* by  $H_2O_2$  was derived from experiments showing that  $\cdot$ OH scavengers (14) decreased killing of high iron *S. aureus* (1.7  $\mu$ g/ $10^9$  organisms) by  $H_2O_2$  (Table I), as well as low iron *S. aureus* (0.2  $\mu$ g/ $10^9$  organisms, data not shown), in close relationship to their effectiveness as  $\cdot$ OH scavengers (Table II). Specifically, increasing concentrations of thiourea, dimethyl thiourea, sodium benzoate, and dimethyl sulfoxide decreased killing of high iron (1.7  $\mu$ g/ $10^9$  organisms) *S. aureus* by  $H_2O_2$  (Table I). In complementary parallel studies, comparable 100 mM concentrations of thiourea, dimethyl thiourea, and sodium benzoate inhibited irradiation-generated  $\cdot$ OH-mediated  $CH_4$  production from  $Me_2SO$  (11) while mannitol, urea, dimethyl urea, and superoxide dismutase had no inhibitory effect (Table II). Thus, the relative effectiveness of these scavengers in inhibiting  $H_2O_2$ -mediated killing of *S. aureus* by  $H_2O_2$  was the same as their relative ability to inhibit

TABLE I

Effect of inhibitors on killing of *S. aureus* by hydrogen peroxide

Inhibitor added	Inhibition of bactericidal activity (% of control with no inhibitor added)
Thiourea (1.5 mM)	7 $\pm$ 3 (18) <sup>a,b</sup>
Thiourea (5.0 mM)	72 $\pm$ 5 (22) <sup>b</sup>
Thiourea (15 mM)	98 $\pm$ 1 (14) <sup>b</sup>
Thiourea (100 mM)	96 $\pm$ 2 (12) <sup>b</sup>
Dimethyl thiourea (2.5 mM)	19 $\pm$ 2 (14) <sup>b</sup>
Dimethyl thiourea (5.0 mM)	35 $\pm$ 2 (14) <sup>b</sup>
Dimethyl thiourea (10 mM)	61 $\pm$ 3 (14) <sup>b</sup>
Dimethyl thiourea (100 mM)	96 $\pm$ 2 (14) <sup>b</sup>
Sodium benzoate (100 mM)	26 $\pm$ 6 (12) <sup>b</sup>
$Me_2SO$ (100 mM)	8 $\pm$ 3 (12) <sup>c</sup>
$Me_2SO$ (140 mM)	19 $\pm$ 2 (11) <sup>b</sup>
$Me_2SO$ (280 mM)	38 $\pm$ 3 (12) <sup>b</sup>
Mannitol (50 mM)	0 (12) <sup>c</sup>
Mannitol (100 mM)	6 $\pm$ 3 (12) <sup>c</sup>
Urea (15 mM)	0 (6) <sup>c</sup>
Urea (100 mM)	6 $\pm$ 1 (12) <sup>c</sup>
Dimethyl urea (100 mM)	0 (14) <sup>c</sup>
Superoxide dismutase (10 $\mu$ g/ml)	2 $\pm$ 1 (16) <sup>c</sup>
Superoxide dismutase (1 $\mu$ g/ml)	0 (18) <sup>c</sup>

<sup>a</sup> Mean  $\pm$  S.E. (number of determinations) from tubes containing *S. aureus* ( $5 \times 10^6$ ),  $H_2O_2$  ( $8.82 \times 10^{-5}$  M), human serum albumin (0.1%), and the inhibitors. In the absence of inhibitors,  $H_2O_2$  killed 52  $\pm$  3.6% of the initial inoculum of *S. aureus* (30 determinations).

<sup>b</sup> Value significantly different ( $p < 0.05$ ) from value with no inhibitor added.

<sup>c</sup> Value not significantly different ( $p > 0.05$ ) from value with no inhibitor added.

TABLE II

Effect of inhibitors on production of  $CH_4$  from  $Me_2SO$  exposed to irradiation

Inhibitor added	Inhibition of $CH_4$ production (% of control with no inhibitor added)
Thiourea (100 mM)	37 $\pm$ 5 (5) <sup>a,b</sup>
Dimethyl thiourea (100 mM)	31 $\pm$ 3 (5) <sup>b</sup>
Sodium benzoate (100 mM)	18 $\pm$ 2 (5) <sup>b</sup>
$Me_2SO$ (100 mM) <sup>c</sup>	
Mannitol (100 mM)	0 (5) <sup>d</sup>
Urea (100 mM)	0 (5) <sup>d</sup>
Dimethyl urea (100 mM)	0 (5) <sup>d</sup>
Superoxide dismutase (100 $\mu$ g)	0 (5) <sup>d</sup>

<sup>a</sup> Mean  $\pm$  S.E. (number of determinations) from tubes containing  $Me_2SO$  (100 mM), human serum albumin (0.1%), and the inhibitor. In the absence of inhibitors, irradiation produced 0.32  $\pm$  0.04 pmol of  $CH_4$  (12 determinations).

<sup>b</sup> Value significantly different ( $p < 0.05$ ) from value with no inhibitor added.

<sup>c</sup> DMSO was not tested against itself.

<sup>d</sup> Value not significantly different ( $p > 0.05$ ) from value with no inhibitor added.

irradiation-generated  $\cdot$ OH-mediated  $CH_4$  production from  $Me_2SO$ .

Additional control observations supported the relative specificity of the inhibitors. First, the active inhibitors did not decrease killing by directly altering viability of *S. aureus*. Specifically, in the absence of  $H_2O_2$ , the numbers of *S. aureus* present at the end of the bactericidal assay were the same with or without the inhibitors (data not shown). Second, under our test conditions, neither thiourea, dimethyl thiourea, sodium benzoate, nor  $Me_2SO$  reacted directly or significantly with  $H_2O_2$  as monitored by optical density at 240 nm (12). In addition, methane, a product of the interaction of  $\cdot$ OH and  $Me_2SO$  (10, 11), was detectable in mixtures of *S. aureus* +  $H_2O_2$  +  $Me_2SO$  or  $FeSO_4$  +  $H_2O_2$  +  $Me_2SO$ , but not detectable in mixtures of  $H_2O_2$  +  $Me_2SO$  or  $FeSO_4$  +  $Me_2SO$  or the above substances individually (data not shown) (10, 11).




FIG. 2. Schematic diagram of the proposed mechanism for killing of bacteria by hydrogen peroxide.  $H_2O_2$  reacts with bacterial iron ( $Fe^{2+}$ ) to form the more toxic hydroxyl radical which damages the bacteria. Scavengers of  $\cdot OH$ , such as dimethyl sulfoxide ( $DMSO$ ), react and inactivate  $\cdot OH$  and protect bacteria from killing by  $H_2O_2$ . When  $Me_2SO$  is the scavenger, methane, a product of the interaction of  $Me_2SO$  and  $\cdot OH$ , is produced.

#### DISCUSSION

Our results indicate that hydroxyl radical may contribute to killing of *S. aureus* by  $H_2O_2$  (Fig. 2). Production of  $\cdot OH$  appeared to involve reaction of  $H_2O_2$  with  $Fe^{2+}$  which had been previously incorporated by *S. aureus*. These statements are corroborated primarily by experiments showing that killing by  $H_2O_2$  was clearly proportional to the intrinsic iron content of the *S. aureus* and that increasing concentrations of  $\cdot OH$  scavengers increasingly inhibited killing of *S. aureus* by  $H_2O_2$ . The additional observation that the degree of inhibition by these agents was proportional to their purported effectiveness as  $\cdot OH$  scavengers further supported this premise. Thus, while iron is an essential growth nutrient, it may also provide an "Achilles heel" for *S. aureus*. Thus, even though *S. aureus* have large amounts of catalase (15), reaction of unscavenged  $H_2O_2$  with  $Fe^{2+}$  could produce  $\cdot OH$  at strategic locations in the bacterium (4-7). Since there are no known enzymatic scavengers of  $\cdot OH$  and presumed nonenzymatic scavengers

may not be fully protective, even small amounts of the highly toxic  $\cdot OH$  produced by this mechanism might be very effective in killing microbial invaders (16-18). Furthermore, one might speculate that this proposed mechanism may contribute to the toxicity of  $H_2O_2$  to other tissues, such as erythrocytes (19), tumor cells (20), or the lung (21).

#### REFERENCES

- Weinberg, E. D. (1974) *Science* **184**, 952-956
- Arbeter, A., Echeverri, L., Franco, D., Munson, D., Velez, H., and Vitale, J. (1971) *Fed. Proc.* **30**, 1421-1430
- Repine, J. E., Berger, E. M., Fox, R. B., and Harada, R. N. (1981) *Infect. Immun.* **32**, 407-410
- McCord, J. M., and Day, E. D., Jr. (1978) *FEBS Lett.* **86**, 139-142
- McCord, J. M. (1974) *Science* **185**, 529-531
- Lai, C. S., and Piette, L. H. (1979) *Tetrahedron Lett.* 775-782
- Walling, C. (1975) *Accts. Chem. Res.* **8**, 125-128
- Carter, P. (1971) *Anal. Biochem.* **40**, 450-451
- Clawson, C. C., and Repine, J. E. (1976) *J. Lab. Clin. Med.* **88**, 316-322
- Repine, J. E., Eaton, J. W., Anders, M. W., Hoidal, J. R., and Fox, R. B. (1979) *J. Clin. Invest.* **64**, 1642-1651
- Repine, J. E., Pfenninger, O. W., Talmage, D. W., Berger, E. M., and Pettijohn, D. E. (1981) *Proc. Natl. Acad. Sci. U. S. A.* **78**, 1001-1003
- Chance, B. (1954) *Methods Biochem. Anal.* **1**, 408-426
- Misra, H. P., and Fridovich, I. (1977) *Arch. Biochem. Biophys.* **183**, 511-519
- Dorfman, L. M., and Adams, G. E. (1978) *NSRDS-Nat. Bur. Stand.* **46**, 9-11
- Mandell, G. L. (1975) *J. Clin. Invest.* **55**, 561-570
- Krinsky, N. I. (1964) *Science* **186**, 363-365
- Fridovich, I. (1978) *Science* **201**, 875-880
- Hassan, H. M., and Fridovich, I. (1979) *Rev. Infect. Dis.* **1**, 357-367
- Weiss, S. J. (1980) *J. Biol. Chem.* **255**, 9912-9917
- Nathan, C. F., Silverstein, S. C., Bruckner, L. H., and Cohn, Z. A. (1979) *J. Exp. Med.* **149**, 100-113
- Shasby, D. M., Van Benthuyzen, K. M., Tate, R. M., McMurtry, I. F., and Repine, J. E. (1981) *Am. Rev. Respir. Dis.* **123**, 243 (abstr.)